

# NEW JERSEY ADULT MOSQUITO SURVEILLANCE

Report start to 1 June 2018, CDC Week 22

Prepared by Lisa M. Reed, Diana Carle and Dina Fonseca  
Center for Vector Biology



This New Jersey Agricultural Experiment Station report is supported by Rutgers University, Hatch funds, funding from the NJ State Mosquito Control Commission and with the participation of the 21 county mosquito control agencies of New Jersey.


**Summary Table – Week 22**

Region	<i>Aedes vexans</i>			<i>Culex Mix</i>			<i>Coquillettidia perturbans</i>			<i>Aedes sollicitans</i>		
	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase
Agricultural	13.52	1.43	4	4.86	0.97	4	0.00	0.04	0	0.00	0.06	0
Coastal	7.81	0.94	4	0.60	1.08	0	0.00	0.03	0	0.71	0.96	0
Delaware Bayshore	0.00	1.89	0	0.00	4.31	0	0.00	0.25	0	0.00	0.94	0
Delaware River Basin	27.86	2.20	4	5.36	0.50	4	0.00	0.10	0	0.14	0.03	4
New York Metro	0.00	1.69	0	0.00	1.91	0	0.00	0.00	0	0.00	0.13	0
North Central Rural	0.04	0.40	0	0.23	0.38	0	0.07	0.00		0.00	0.00	0
Northwest Rural	0.23	1.59	0	0.37	0.73	0	0.00	<0.01	0	0.00	0.00	0
Philadelphia Metro	0.00	3.30	0	0.00	2.13	0	0.00	0.04	0	0.00	0.00	0
Pinelands	6.40	0.48	4	0.14	0.75	0	0.00	0.13	0	0.00	0.00	0
Suburban Corridor	0.00	2.37	0	0.06	0.94	0	0.03	0.17	0	0.00	<0.01	0

\*Averages represent data from, at most, the previous 5 years. Increase is a scale of current values from historical values where no difference or a decrease is represented by 0 (blue), up to 50% greater difference by 1 (green), up to 100% greater difference by 2 (yellow), up to 150% greater difference by 3 (orange) and greater than 150% increase by 4 (red). White cells in the increase column denote increases from an historic zero and thus no value can be appropriately given. nd=no data reported.


State Summary: Our mosquito season in New Jersey had started many weeks ago with cold-tolerant species on the wing during really variable weather (read short-sleeves hot and then snowy cold) as spring developed. Now that the snow is behind us, the mosquito season is in full force. Recent significant rainfall has produced *Aedes vexans* populations in amounts significantly above historical levels in the Agricultural, Coastal, Delaware River Basin and Pinelands regions. Also elevated are *Culex Mix* in the Agricultural and Delaware River Basin regions and *Aedes sollicitans* in the Delaware River Basin. Noteworthy is the presence of *Coquillettidia perturbans* in the North Central Rural regions, when it has not been recorded in the traps there this early in the season.

## Climate Factors




The three figures show the interpolation of average maximum (°F) and minimum temperature (°F) and total precipitation (inches) for 30 days prior to 1 June 2018 in New Jersey. Data points are from about 54 weather stations maintained through the New Jersey Weather & Climate Network and the State Climatologist. Interpolation between points was performed using ArcMap 10.1.

**The Species Graphs:** The species graph pages include a graph with two plots for each of the ten regions defined on the first page (Agricultural, Coastal, Delaware Bayshore, Delaware River, New York Metro, North-Central, Northwestern, Philadelphia Metro, Pinelands, and Suburban Corridor). Below is an example of one graph from one species within one region. The bar plot show the average number of mosquitoes per trap within the region (weekly means) and line plots show the historical trend as the average number of mosquitoes from the previous 5 years (5-year average). In general, historical data are running means from the previous 5 years, but on occasion, will include data from fewer years. Adjustments are made to account for year discrepancies. Data for this week are from Atlantic, Essex, Hunterdon, Morris, Passaic, Salem, and Warren counties. Data for the previous week are from Atlantic, Bergen, Burlington, Cape May, Cumberland, Essex, Hunterdon, Middlesex, Monmouth, Morris, Ocean, Passaic, Salem, Sussex and Warren counties.




# Aedes vexans - Fresh Floodwater Species

## Multivoltine Aedine (Ae. vexans Type)




# Culex Mix – Permanent Water Species

## Multivoltine *Culex/Anopheles* (Cx. *pipiens* Type)




# *Culiseta melanura* – Miscellaneous Group

## Unique (*Cs. melanura* Type)



# *Aedes sollicitans* - Salt Floodwater Species

## Multivoltine Aedine (*Ae. sollicitans* Type)



# *Coquillettidia perturbans*

## Monotypic (*Coquillettidia perturbans* Type)


Agricultural	Coastal	Delaware Bayshore	Delaware River Basin
New York Metro	North Central Rural	Northwestern Rural	Philadelphia Metro
Pinelands	Suburban Corridor	Comments	
		<p><i>Coquillettidia perturbans</i> populations are located in cattail swamps and other wetlands with emergent vegetation, often making control of their numbers dependent on controlling water depth. This mid-season species is a potential inland vector of EEE and should be monitored closely when the arbovirus is present. Noteworthy (although difficult to see on this graph) is the presence of this species when the previous 5-year records do not report it this early in the North Central Rural region.</p>	

WNV

EEE


**Top Ten Mosquito Species/Region** - ■ Ae. albopictus, ■ Ae. japonicus (invasives); ■ Cs. melanura or Cx. erraticus ■ Coq. perturbans

Note: In early season when fewer species are caught, graphs may show less than ten species/region or 25 statewide.




## Agricultural

Total # mosquitoes




## Coastal

Total # mosquitoes




### Delaware Bayshore

#### Total # mosquitoes




### Delaware River Basin

#### Total # mosquitoes




## New York Metropolitan

### Total # mosquitoes




## North Central Rural

### Total # mosquitoes




## Northwest Rural

Total # mosquitoes




## Philadelphia Metropolitan

Total # mosquitoes



## Pinelands

Total # mosquitoes



## Suburban Corridor

Total # mosquitoes

