

NEW JERSEY ADULT MOSQUITO SURVEILLANCE Report

14 July to 20 July 2019, CDC Week 29

Prepared by Lisa M. Reed and Dina Fonseca

Center for Vector Biology



This New Jersey Agricultural Experiment Station report is supported by Rutgers University, Hatch funds, funding from the NJ State Mosquito Control Commission and with the participation of the 21 county mosquito control agencies of New Jersey.

Summary Table –begin to Week 29

Region	<i>Aedes vexans</i>			<i>Culex Mix</i>			<i>Coquillettidia perturbans</i>			<i>Aedes sollicitans</i>		
	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase
Agricultural	1.38	4.25	0	4.98	5.60	0	0.10	1.06	0	0.07	0.19	0
Coastal	0.71	4.67	0	9.56	7.07	1	0.95	0.53	2	7.76	5.34	1
Delaware Bayshore	0.79	1.03	0	5.10	20.08	0	1.03	1.35	0	1.71	1.87	0
Delaware River Basin	3.64	17.11	0	16.93	4.84	4	1.50	2.33	0	0.00	0.00	0
New York Metro	0.39	7.08	0	3.30	8.69	0	0.00	0.15	0	0.00	0.89	0
North Central Rural	0.02	0.32	0	0.14	0.45	0	0.00	0.47	0	0.00	0.00	0
Northwest Rural	1.26	10.89	0	1.74	5.07	0	3.60	1.50	3	0.00	0.00	0
Philadelphia Metro	1.14	6.10	0	1.00	4.40	0	2.86	0.22	4	0.00	0.00	0
Pinelands	0.21	1.37	0	1.25	3.31	0	0.91	1.84	0	0.79	0.11	4
Suburban Corridor	1.10	2.59	0	0.83	1.31	0	0.13	0.55	0	0.00	0.00	0

*Averages represent data from, at most, the previous 5 years. Increase is a scale of current values from historical values where no difference or a decrease is represented by 0 (blue), up to 50% greater difference by 1 (green), up to 100% greater difference by 2 (yellow), up to 150% greater difference by 3 (orange) and greater than 150% increase by 4 (red). White cells in the increase column denote increases from an historic zero and thus no value can be appropriately given. nd=no data reported.

State Summary: While *Aedes vexans* populations remain below historical averages, other populations show recent increases. *Culex Mix* populations were again significantly higher in the Delaware River Basin and mildly higher in the Coastal region. *Coquillettidia perturbans* was significantly higher in the Philadelphia Metropolitan region, moderately higher in the Northwest Rural and the Coastal regions. *Aedes sollicitans* abundances were significantly above historical averages in the Pinelands and mildly elevated in the Coastal region.

Climate Factors


Average High Temperature	Average Low Temperature	Total Precipitation
Not available	Not available	Not available

The three figures show the interpolation of average maximum (°F) and minimum temperature (°F) and total precipitation (inches) for 30 days prior to 20 July 2019 in New Jersey. Data points are from about 50 weather stations maintained through the New Jersey Weather & Climate Network and the State Climatologist. Interpolation between points was performed using ArcMap 10.1.




The Species Graphs: The species graph pages include a graph with two plots for each of the ten regions defined on the first page (Agricultural, Coastal, Delaware Bayshore, Delaware River, New York Metro, North-Central, Northwestern, Philadelphia Metro, Pinelands, and Suburban Corridor). Below is an example of one graph from one species within one region. The bar plot shows the average number of mosquitoes per trap within the region (weekly means) and line plots show the historical trend as the average number of mosquitoes from the previous 5 years (5-year average). In general, historical data are running means from the previous 5 years, but on occasion, will include data from fewer years. Adjustments are made to account for year discrepancies. Data for this week are from Atlantic, Burlington, Cape May, Cumberland, Hudson, Hunterdon, Mercer, Monmouth, Passaic, Salem, Somerset, Sussex, Union, and Warren counties. Data for the previous week are from Atlantic, Burlington, Cape May, Cumberland, Hudson, Hunterdon, Mercer, Monmouth, Passaic, Salem, Somerset, Sussex, Union, and Warren counties.

Weekly Means Against 5-year Average




Aedes vexans - Fresh Floodwater Species


Multivoltine Aedine (Ae. vexans Type)




Culex Mix – Permanent Water Species Multivoltine *Culex/Anopheles* (*Cx. pipiens* Type)



Culiseta melanura – Miscellaneous Group Unique (Cs. melanura Type)




Aedes sollicitans - Salt Floodwater Species Multivoltine Aedine (Ae. sollicitans Type)



Coquillettidia perturbans

Monotypic (Coquillettidia perturbans Type)




WNV

EEE


Top Ten Mosquito Species/Region - ■ *Ae. albopictus*, ■ *Ae. japonicus* (invasives); ■ *Cs. melanura* or *Cx. erraticus* ■ *Coq. perturbans*

Note: In early season when fewer species are caught, graphs may show less than ten species/region or 25 statewide.




Agricultural

Total # mosquitoes




Coastal

Total # mosquitoes




Delaware Bayshore

Total # mosquitoes




Delaware River Basin

Total # mosquitoes




New York Metropolitan

Total # mosquitoes




North Central Rural

Total # mosquitoes




Northwest Rural

Total # mosquitoes




Philadelphia Metropolitan

Total # mosquitoes



Pinelands



Suburban Corridor

